

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁴ : C12N 15/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 89/ 02468 (43) International Publication Date: 23 March 1989 (23.03.89)</p>																		
<p>(21) International Application Number: PCT/US88/03089 (22) International Filing Date: 8 September 1988 (08.09.88) (31) Priority Application Number: 096,074 (32) Priority Date: 11 September 1987 (11.09.87) (33) Priority Country: US (71) Applicants: WHITEHEAD INSTITUTE FOR BIOMEDICAL RESEARCH [US/US]; Nine Cambridge Center, Cambridge, MA 02142 (US). HOWARD HUGUES MEDICAL INSTITUTE [US/US]; 6701 Rockledge Drive, Bethesda, MA 20817 (US). (72) Inventors: MULLIGAN, Richard, C. ; 441 Franklin Street, Cambridge, MA 02138 (US). WILSON, James, M. ; 24 Whitman Road, Waltham, MA 02154 (US).</p>		<p>(74) Agents: GRANAHAN, Patricia et al.; Hamilton, Brook, Smith & Reynolds, Two Militia Drive, Lexington, MA 02173 (US). (81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>																		
<p>(54) Title: TRANSDUCED FIBROBLASTS AND USES THEREFOR</p> <div data-bbox="386 1150 1208 1871"> <table border="1"> <caption>Data points estimated from the graph</caption> <thead> <tr> <th>Time (Hours)</th> <th>FWR 2ng/hr./10⁶ (ng/10⁶ cells)</th> <th>NIH 3T3 0.4ng/hr./10⁶ (ng/10⁶ cells)</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>0</td> <td>0</td> </tr> <tr> <td>6</td> <td>11</td> <td>2</td> </tr> <tr> <td>12</td> <td>24</td> <td>4</td> </tr> <tr> <td>24</td> <td>46</td> <td>10</td> </tr> <tr> <td>36</td> <td>38</td> <td>10</td> </tr> </tbody> </table> </div> <p>(57) Abstract</p> <p>Fibroblasts transduced with genetic material encoding a polypeptide or protein of interest and, optionally, a selectable marker, as well as methods for making and using the transduced fibroblasts. Such fibroblasts are useful in delivering the encoded polypeptide or protein, such as an enzyme, a hormone or a drug, to an individual who has had a graft or implant of the transduced cells.</p>			Time (Hours)	FWR 2ng/hr./10 ⁶ (ng/10 ⁶ cells)	NIH 3T3 0.4ng/hr./10 ⁶ (ng/10 ⁶ cells)	0	0	0	6	11	2	12	24	4	24	46	10	36	38	10
Time (Hours)	FWR 2ng/hr./10 ⁶ (ng/10 ⁶ cells)	NIH 3T3 0.4ng/hr./10 ⁶ (ng/10 ⁶ cells)																		
0	0	0																		
6	11	2																		
12	24	4																		
24	46	10																		
36	38	10																		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

-1-

TRANSDUCED FIBROBLASTS AND USES THEREFORDescriptionBackground

05 In the embryo, the mesoderm is the middle of the
three primary third germ layers, and lies between the
ectoderm and the entoderm. It gives rise, during
development, to the connective tissues, all body
musculature, blood, cardiovascular and lymphatic
10 systems, most of the urogenital system and the lining
of the pericardial, pleural and peritoneal cavities.
That portion of the mesoderm which produces the
connective tissue, blood vessels and blood, the
lymphatic system and the heart is referred to as the
mesenchyme. One type of cells produced by
15 mesenchymal cells is fibroblast cells, which are
stellate or spindle-shaped cells with cytoplasmic
processes. They are present in connective tissue and
are capable of forming collagen fibers.

20 Fibroblasts, like other cells in the body,
contain an entire complement of all genetic material.
However, only a small percentage of the genes they
contain are expressed at biologically functional
levels. That is, most of the genes in fibroblasts
are not expressed at all, or are expressed at such
25 low levels that the polypeptides they encode are

-2-

produced in undetectable amounts or concentrations which are biologically nonfunctional or insignificant.

- It is possible, using methods developed in recent years, to attain interspecies genetic recombination. Genes derived from different biological classes are able to replicate and be expressed in a selected microorganism. Therefore, it is possible to introduce into a microorganism genes specifying a metabolic or synthetic function (e.g., hormone synthesis, protein synthesis, nitrogen fixation) which is characteristic of other classes of organisms by linking the genes to a particular viral or plasmid replicon.
- Since the late 1970s, progress has been made toward the development of general methods for introducing cloned DNA sequences into mammalian cells. At the present time, however, there is a need for an effective method of stably introducing genetic material into fibroblasts and enabling them to express genetic material which they do not normally express, or normally express at biologically insignificant levels.

Summary of the Invention

- The invention described herein relates to genetically engineered mesenchymal or connective tissue cells and particularly to genetically engineered fibroblasts which express in biologically significant concentrations genetic material which is not normally expressed in significant concentrations in such cells. It also relates to methods of stably

-3-

introducing into fibroblasts such genetic material and methods of using the genetically engineered fibroblasts.

Fibroblasts of this invention have stably
5 incorporated in them genetic material of interest and express the incorporated genetic material. This genetic material of interest is referred to herein as incorporated genetic material. The incorporated genetic material can be DNA or RNA which does not
10 normally occur in fibroblasts; DNA or RNA which normally occurs in fibroblasts but is not expressed in them at levels which are biologically significant (i.e., levels sufficient to produce the normal physiological effects of the polypeptide it encodes);
15 DNA or RNA which occurs in fibroblasts and has been modified so that it is expressed in fibroblasts; and any DNA or RNA which can be modified to be expressed in fibroblasts, alone or in any combination thereof. The incorporated genetic material expressed by
20 fibroblasts of the present invention can include genetic material encoding a selectable marker, thus providing a means by which cells expressing the incorporated genetic material are identified and selected for. Fibroblasts containing incorporated
25 genetic material are referred to as transduced fibroblasts.

In particular, retroviral vectors have been used to stably transduce fibroblasts with genetic material which includes genetic material encoding a polypep-
30 tide or protein of interest not normally expressed at biologically significant levels in fibroblasts. The genetic material introduced in this manner also

-4-

included genetic material encoding a dominant selectable marker. Genetic material including DNA encoding a polypeptide of interest and DNA encoding a dominant selectable marker has been introduced into
5 cultured fibroblasts. Expression of these genes by the fibroblasts into which they have been incorporated (i.e., fibroblasts transduced by the use of retroviral vectors) has also been demonstrated.

A method of transplanting transduced fibroblasts
10 which express the incorporated genetic material they contain is also a subject of the present invention. Transduced fibroblasts of the present invention are used for the constitutive delivery of polypeptides or proteins, useful in prevention and therapy or
15 treatment, which are presently administered parenterally. They can be used in skin grafts and in glial cell or fibroblast implants which introduce DNA encoding a polypeptide or protein of interest into the central nervous system.

20 There are many advantages to fibroblasts of the present invention and the delivery system for a polypeptide or protein of interest which make them very useful. For example, a skin graft using fibroblasts which include incorporated genetic
25 material encoding a polypeptide of interest (e.g., a hormone, enzyme, drug) synthesizes the encoded polypeptide and thus serves as a continuous delivery system for that polypeptide. In this way, the hormone or other polypeptide is diffused into the
30 bloodstream of the individual receiving the skin graft.

-5-

An important advantage is that the genetically engineered fibroblasts of the present invention can be used to administer therapeutic proteins (e.g., hormones, enzymes, clotting factors) which are presently administered intravenously, intramuscularly or subcutaneously. In addition, there is no need for extensive (and often costly) purification of the polypeptide before it is administered to an individual, as is generally necessary with an isolated polypeptide (e.g., insulin). Fibroblasts modified according to the present invention produce the polypeptide hormone as it would normally be produced. In the case of insulin, for example, the genetically engineered fibroblasts produce insulin in the same form as that made in the pancreas.

Another advantage to the use of a graft having fibroblasts of the present invention is that by controlling the size of the graft, the amount of the polypeptide delivered to the body can be controlled. In addition, in the case of a skin graft, it can be excised if there is no longer a need for the polypeptide being produced. For example, if delivery of the polypeptide (hormone, enzyme, or drug) is necessary only for a specific period, the engineered graft can be removed when treatment is no longer needed.

Another important advantage of the delivery system of this invention is that because it is a continuous delivery system, the fact that polypeptide hormones have very short half lives is not a limitation. For example, the half life of human growth hormone (HGH) is approximately 19 minutes, of

-6-

parathyroid hormone, approximately 2½ to 5 minutes and, of native insulin (pure insulin), approximately 3 to 4 minutes.

Because genes can be introduced into fibroblasts using a retroviral vector, they can be "on" (subject to) the retroviral vector control; in such a case, the gene of interest is transcribed from a retroviral promoter. A promoter is a specific nucleotide sequence recognized by RNA polymerase molecules that start RNA synthesis. Alternatively, retroviral vectors having additional promoter elements (in addition to the promoter incorporated in the recombinant retrovirus) which are responsible for the transcription of the genetic material of interest, can be used. For example, a construct in which there is an additional promoter modulated by an external factor or cue can be used, making it possible to control the level of polypeptide being produced by the fibroblasts by activating that external factor or cue. For example, heat shock proteins are proteins encoded by genes in which the promoter is regulated by temperature. The promoter of the gene which encodes the metal-containing protein metallothioneine is responsive to cadmium (Cd^{++}) ions. Incorporation of this promoter or another promoter influenced by external cues also makes it possible to regulate the production of the polypeptide by the engineered fibroblasts.

Brief Description of the Drawings

Figure 1 is a schematic representation of a wild type murine leukemia virus (retroviral) genome.

-7-

Figure 2 is a schematic representation of retroviral vectors, each having a recombinant genome, useful in the present invention. Figure 2a is pZIPNeo; Figure 2b is pLJ; Figure 2c is pWe; Figure 2d is pEm; and Figure 2e is pIp.

Figure 3 is a schematic representation of the construction of a recombinant retroviral vector, using the pLJ vector represented in Figure 2b and the human parathyroid hormone gene.

Figure 4 is a graph representing the quantity of human parathyroid hormone produced, in the designated time period, by transduced fibroblasts. The dashed line represents production by transduced NIH 3T3 cells on tissue culture plates. The solid line represents production by transduced fibroblasts derived from a rat and grown on tissue culture plates.

Figure 5 is a graph representing the quantity of human parathyroid hormone produced, in the designated time periods, in rats injected with either transduced fibroblasts alone (blackened circles) or transduced fibroblasts coated onto cyto 3 beads and grown to confluence.

Detailed Description of the Invention

Genetic material of interest has been incorporated into fibroblasts and expressed in the resulting genetically engineered fibroblasts. Genetic material incorporated into fibroblasts according to the method described can be DNA or RNA which does not normally occur in fibroblasts; DNA or RNA which normally occurs in fibroblasts but is not

-8-

expressed in them at levels which are biologically significant (levels sufficient to produce the normal physiological effects of the polypeptide it encodes); DNA or RNA which occurs in fibroblasts and has been
5 modified so that it is expressed in such cells; and any DNA or RNA which can be modified to be expressed in fibroblasts, alone or in a combination thereof. This genetic material of interest is referred to herein as incorporated genetic material. Fibroblasts
10 of the present invention express the incorporated genetic material. The incorporated genetic material (i.e., DNA or RNA) expressed by fibroblasts of the present invention is genetic material encoding a polypeptide or a protein of interest (genetic
15 material of interest), alone or in combination with a gene encoding a selectable marker.

For example, genetic material encoding a hormone has been introduced into fibroblasts by exposing them to media that contains a virus having a recombinant
20 genome (i.e., by infecting them). The media used is obtained by harvesting media in which producer cells (e.g., a Psi am or amphotropic producer) have been grown. That is, producer cells have been grown in tissue culture to a confluent density in Dulbecco's
25 Modified Eagle's medium (DME) with 10% calf serum (CS) and penicillin and streptomycin. Fresh media is added and subsequently (e.g., approximately 12 hours later), the media is harvested. Approximately 10 ml of media is harvested from a 10 cm plate of confluent
30 producer cells. The spent media (or viral stock) is filtered through a 0.45 micron Millipore filter to remove detached producer cells and is used

immediately to infect cells or is stored at -70° C.
Media is removed from a subconfluent plate of
fibroblasts (recipient fibroblasts) and quickly
replaced with viral stock (e.g., 5 ml/10 cm. plate)
5 containing 8 mcg/ml. of Polybrene (Aldrich).
Subsequently (e.g., approximately 12 hours later),
this is removed and replaced with fresh media. Thus,
the media used is a viral supernatant. The
recombinant genome of the infectious virus includes
10 the genetic material of interest. The recombinant
genome can also have genetic material encoding a
dominant selectable marker.

Thus, fibroblasts are made which express a
polypeptide not normally expressed by such cells at
15 biologically significant levels and, optionally, a
dominant selectable marker.

In particular, fibroblasts are exposed to media
containing infectious virus produced in Psi am cells;
the infectious virus contain a recombinant genome
20 having the genetic material of interest. The
recombinant genome in one instance includes genetic
material encoding human parathyroid hormone (hPTH).
Optionally, it can also include a gene encoding a
dominant selectable marker (e.g., the neo gene which
25 encodes neomycin resistance). As a result, the
fibroblasts are transduced -- that is, the genetic
material of interest (in this case, DNA encoding hPTH
and, optionally, the neo gene) is stably introduced
into the fibroblasts. The transduced fibroblasts
30 express the encoded hPTH and, if the neo gene is
present, express it, resulting in cells having the
selectable trait.

-10-

Alternatively, fibroblasts are transduced as a result of exposure to media containing infectious virus in which the recombinant genome includes DNA encoding one or more of the following:

- 5 the receptor for low density lipoproteins;
- human growth hormone;
- the gene that confers resistance to histidonal;
- human adenosine deaminase;
- the receptor for interleukin 2;
- 10 human beta-globin;
- human alpha-globin;
- a mutant form of dihydrofolate reductase;
- multidrug resistance;
- glucose cerebrosidase from humans;
- 15 the E1A gene from adenovirus;
- many different genes for HLA in humans;
- human albumin;
- human ornithine transcarbamylase;
- beta-galactosidase from E. coli;
- 20 resistance to neomycin in E. coli;
- human insulin; and
- the envelope protein from Moloney murine leukemia virus.

Fibroblasts expressing the incorporated genetic

- 25 material are grown to confluence in tissue culture vessels; removed from the culture vessel in which they were grown; and introduced into or applied to the body. They have been introduced into the intraperitoneal cavity, both alone and coated onto
- 30 microcarrier beads (approximately 100 cells/bead)

-11-

having collagen-covered surfaces. Alternatively, they can be applied as a component of a skin graft, using a method such as that described by Bell in U.S. 4,485,096, the teachings of which are incorporated
5 herein by reference.

Fibroblasts expressing the incorporated genetic material can also be introduced into the central nervous system. This can be done, for example, by introducing genetically engineered fibroblasts of the
10 present invention (i.e., fibroblasts into which a gene of interest has been introduced) directly into specific regions of the brain through stereotactic administration. It can also be introduced into the cerebrospinal fluid via lumbar puncture or directly
15 into the ventricles, which would result in seeding of the fibroblasts along the meninges.

Once introduced into or applied to the body, the transduced fibroblasts provide a continuous supply of the hormone, enzyme or drug encoded by the genetic
20 material of interest. In the example described, the encoded product is hPTH.

The amount of the hormone, enzyme or drug supplied in this way can be modified or regulated as needed. This is done, for example, by using external
25 cues or factors which affect their production; by controlling the size of the graft applied or the quantity of fibroblasts introduced into the body; or by removing the graft.

Cultured Fibroblasts

30 Fibroblasts are obtained from a subject by skin biopsy (e.g., a small punch biopsy from any area of

WO 89/02468

PCT/US88/03089

-12-

the body). The resulting tissue is placed in tissue culture media and separated into small pieces (e.g., by use of scalpels to tease apart the tissue). Small chunks of the tissue are placed on a wet surface of a tissue culture flask; approximately 10 pieces are placed in each flask. The flask is turned upside-down, closed tight and left at room temperature overnight. After 24 hours at room temperature, the flask is inverted; the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media with 10% FBS and penicillin and streptomycin) is added. This is then incubated at 37° for approximately one week. At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerges. The monolayer is trypsinized and scaled up into larger flasks. The fibroblasts can be maintained in culture for approximately 50 generations; at approximately that time, they undergo what is called a crisis and subsequently do not grow very well. Soon after the fibroblasts are scaled up to (replated onto) larger flasks, they are infected according to the protocol described below.

25 Retroviral Vectors

Retroviruses are RNA viruses; that is, the viral genome is RNA. This genomic RNA is, however, reverse transcribed into a DNA intermediate which is integrated very efficiently into the chromosomal DNA of infected cells. This integrated DNA intermediate is referred to as a provirus. As shown in Figure 1,

-13-

the retroviral genome and the proviral DNA have three genes: the gag, the pol and the env, which are flanked by two long terminal repeat (LTR) sequences.

The gag gene encodes the internal structural (nucleocapsid) proteins; the pol gene encodes the RNA-directed DNA polymerase (reverse transcriptase); and the env gene encodes viral envelope glycoproteins. The 5' and 3' LTRs serve to promote transcription and polyadenylation of virion RNAs.

10 Adjacent to the 5' LTR are sequences necessary for reverse transcription of the genome (the tRNA primer binding site) and for efficient encapsidation of viral RNA into particles (the Psi site).
Mulligan, R.C., In: Experimental Manipulation of Gene
15 Expression, M. Inouye (ed), 155-173 (1983); Mann, R., et al., Cell, 33:153-159 (1983); Cone, R.D. and R.C. Mulligan, Proceedings of the National Academy of Sciences, U.S.A., 81:6349-6353 (1984).

If the sequences necessary for encapsidation (or
20 packaging of retroviral RNA into infectious virions) are missing from the viral genome, the result is a cis defect which prevents encapsidation of genomic RNA. However, the resulting mutant is still capable of directing the synthesis of all virion proteins.

25 Mulligan and co-workers have described retroviral genomes from which these Psi sequences have been deleted, as well as cell lines containing the mutant stably integrated into the chromosome. Mulligan, R.C., In: Experimental Manipulation of Gene
30 Expression, M. Inouye (ed), 155-173 (1983); Mann, R., et al., Cell, 33:153-159 (1983); Cone, R.D. and R.C. Mulligan, Proceedings of the National Academy of

WO 89/02468

PCT/US88/03089

-14-

Sciences, U.S.A., 81:6349-6353 (1984). The teachings of these publications are incorporated herein by reference.

5 The Psi 2 cell line described by Mulligan and co-workers was created by transfecting NIH 3T3 fibroblasts with pMOV-Psi⁻, which is an ecotropic Moloney murine leukemia virus (Mo-MuLV) clone. pMOV-Psi⁻ expresses all the viral gene products but lacks the Psi sequence, which is necessary for encapsidation of
10 the viral genome. pMOV-Psi⁻ expresses an ecotropic viral envelope glycoprotein which recognizes a receptor present only on mouse (and closely related rodent) cells.

Another cell line is the Psi am line, which are
15 Psi-2-like packaging cell lines. These Psi-am cell lines contain a modified pMOV-Psi-genome, in which the ecotropic envelope glycoprotein has been replaced with envelope sequences derived from the amphotropic virus 4070A. Hartley, J.W. and W.P. Rowe, Journal of
20 Virology, 19:19-25 (1976). As a result, they are useful for production of recombinant virus with amphotropic host range. The retrovirus used to make the Psi am cell line has a very broad mammalian host range (an amphotropic host range) and can be used to
25 infect human cells. If the recombinant genome has the Psi packaging sequence, the Psi-am cell line is capable of packaging recombinant retroviral genomes into infectious retroviral particles. Cone, R. and Mulligan, R., Proceedings of the National Academy of
30 Sciences, USA, 81:6349-6353 (1984).

The retroviral genome has been modified by Cone and Mulligan for use as a vector capable of

-15-

introducing new genes into cells. As shown in Figure 2, the gag, the pol and the env genes have all been removed and a DNA segment encoding the neo gene has been inserted in their place. The neo gene serves as a dominant selectable marker. The retroviral sequence which remains part of the recombinant genome includes the LTRs, the tRNA binding site and the Psi packaging site. Cepko, C. et al., Cell, 37:1053-1062 (1984).

10 Additional vector constructions which have been used in producing transduced fibroblasts of the present invention are represented in Figure 2 and are described in detail below.

pZip. The construction of this vector has been described in Cepko, C.L. et al. Cell 37:1053 (1984). Briefly, this vector is capable of expressing two genes: the gene of interest and the Neo gene as a selectable marker.

The gene of interest is cloned into a BamHI site just distal to the 5' LTR, flanked by a splice donor site, and a splice acceptor site. Two transcripts will result from transcription of the provirus: the unprocessed transcript will result in expression of the gene of interest while the "processed" transcript will result in expression of the Neo gene.

pLJ. The characteristics of this vector have been described in Korman, A.J. et al., Proceedings of the National Academy of Sciences, USA 84:2150 (1987). This vector is capable of expressing two genes: the gene of interest and a dominant selectable marker, such as the Neo gene. The gene of interest is cloned in direct orientation into a BamHI/SmaI/SalI cloning

-16-

site just distal to the 5' LTR, while, the Neo gene is placed distal to an internal promoter (from SV40) which is farther 3' than is the cloning site (is located 3' of the cloning site). Transcription from PLJ is initiated at two sites: 1) the 5' LTR, which is responsible for expression of the gene of interest and 2) the internal SV40 promoter, which is responsible for expression of the Neo gene.

pWe. The construction and initial characterization of this vector have been described. Choudory, P.V. et al, CSH Symposia Quantitative Biology, L.I. 1047 (1986). Briefly, this vector can drive expression of two genes: a dominant selectable marker, such as Neo, which is just downstream from the 5' LTR and a gene of interest which can be cloned into a BAMH1 site just downstream from an internal promoter capable of high level constitutive expression. Several different internal promoters, such as the beta-actin promoter from chicken (Choudory, P.V. et al, CSH Symposia Quantitative Biology, L.I. 1047 (1986)), and the histone H4 promoter from human (Hanly, S.M. et al., Molecular and Cellular Biology 5:380 (1985)) have been used. Expression of the Neo gene is from a transcript initiated at the 5' LTR; expression of the gene of interest is from a transcript initiated at the internal promoter.

pEm. In this simple vector, the entire coding sequence for gag, pol and env of the wild type virus is replaced with the gene of interest, which is the only gene expressed. The components of the pEm vector are described below. The 5' flanking

-17-

sequence, 5' LTR and 400 bp of contiguous sequence (up to the BAMHI site) is from pZIP. The 3' flanking sequence and LTR are also from pZIP; however, the cla site 150 bp upstream from the 3' LTR has been

5 linker with BAMHI and forms the other half of the BamHI cloning site present in the vector. The HindIII/EcoRI fragment of pBR322 forms the plasmid backbone. This vector is derived from sequences cloned from a strain of Moloney Murine Leukemia

10 virus. An analogous vector has been constructed from sequences derived from the myeloproliferative sarcoma virus.

pIp. This vector is capable of expressing a single gene driven from an internal promoter. The

15 construction of these vectors is summarized below. The 5' section of the vector, including the 5' flanking sequences, 5' LTR, and 1400 bp of contiguous sequence (up to the xho site in the gag region) is derived from wild type Moloney Leukemia virus

20 sequence. Shinnick et al., Nature, 293:543 (1918). The difference between the two is that a SacII linker is cloned into an HaeIII restriction site immediately adjacent to the ATG of the gag gene. The 3' section of the vector, including the 3' flanking sequences,

25 3' LTR and 3' contiguous sequence (up to the clal site in the env coding region) is from pZIP. However, there are two modifications: 1) the clal site has been linked to BamHI and 2) a small sequence in the 3' LTR spanning the enhancer (from PvuII to

30 XbaI) has been deleted. Bridging the 5' and 3' sections of the vector is one of several promoters; each one is contained on a xhoI/BamHI fragment, and

-18-

each is capable of high level constitutive expression in most tissues. These promoters include beta-actin from chicken (Choudory, P.V. et al, CSH Symposia Quantitative Biology, L.I. 1047 (1986), and
5 thymidine Kinase from Herpes Simplex Virus, histone H4 from human (Hanly, S.M. et al., Molecular and Cellular Biology 5:380 (1985)). The vector backbone is the HindIII/EcoRI fragment from pBR322. The gene of interest is cloned into the BamHI site in direct
10 orientation, just downstream from the internal promoter.

RO vectors. This category represents a heterogeneous group of vectors in which the gene of interest contains all the sequences necessary for
15 transcription (i.e., promoter/enhancer, coding sequence with and without introns, and polyadenylation signal) and is introduced into the retroviral vector in an orientation in which its transcription is in a direction opposite to that of
20 normal retroviral transcription. This makes it possible to include more of the cis-acting elements involved in the regulation of the introduced gene. Virtually, any of the above described genes can be adapted to be a RO vector. One example is described
25 by Cone et al. in which the entire beta globin gene was cloned in reverse orientation into the BamHI site of pZip. Cone, R. and R.C. Mulligan Proceeding of the National Academy of Sciences, U.S.A.,
81:6349-6353 (1984). RO vectors have been
30 constructed in which the gene of interest has been cloned into the XhoI/BamHI site of pIp (essentially replacing the internal promoter).

-19-

Vectors into which a gene of interest has been inserted are shown in Table 1 as well as results obtained when the construction was tested for expression.

-20-

Table 1 Vectors Containing Gene of Interest

TABLE 1

<u>Protein of Function</u>		<u>Vector</u>					
		pZip	pLJ	pWe	pEm	PIP	pRO
5	Human growth hormone	3	3				3
	Human PTH	3	3		3	3	
	Human receptor for LDL				3	3	
	Human albumin		3	2			1
	Human ornithine transcarbamylase				1	1	
10	Human adenosine deaminase					3	
	Receptor for interleukin-2			3	3		
	Human Beta globin						3
	Human alpha globin						3
	Mutant dihydrofolate reductase	3			3	3	
15	Multidrug resistance					3	
	Human glucose cerebrosidase			3			
	Neomycin	3	3	3	3	3	
	EIA gene from aderovirus	3					
	Histidinol ^R from <u>E. coli</u>	3					
20	-galactosidase from <u>E. coli</u>		3				
	HLA antigens from human		3				
	Human insulin		3				
	Envelope from Moloney MLV	1		3			
1	constructed, not tested for expression						
25	2	constructed, no expression when tested					
	3	constructed, expression detected					

-21-

Vectors which include a selectable marker are represented in Figure 2a, 2b and 2c. In Figure 2a, pZIP neo SVX, described by Cepko is represented. Cepko, C. et al., Cell, 37:1053-1062 (1984). Here, there is a splice donor and splice accept, noted as SD and SA, between which the gene of interest is inserted, using known techniques. Expression of the inserted gene is based on the LTR in the unprocessed message; the processed message is responsible for expressing neomycin.

In Figure 2b, vector pLJ is represented. In pLJ, the genetic material of interest is inserted just following the 5' LTR. Expression of this genetic material is transcribed from the LTR and expression of the neo gene is transcribed from an internal SV40 promoter.

In Figure 2c, the pWe vectors are represented. In pWe vectors, the LTR promoter and transcript is responsible for expressing neomycin and an internal promoter is responsible for expression of the gene of interest. Promoters useful in this type of vector include promoters derived from chicken beta-actin, human histone, Herpes simplex thymidine kinase, thy 1 (a tissue-specific promoter used in T cells) and rat albumin.

Vectors without a selectable marker can also be used to transduce fibroblasts with genetic material of interest. Such vectors are basically simplifications of the vectors previously described, in which there is such a marker. Vector pEm is represented in Figure 2d; as represented, the main components of the vector are the 5' and 3' LTR, and the genetic

-22-

material of interest, inserted between the two LTRs. pIp represents a series of useful vectors in which there is an internal promoter. As represented in Figure 2e, there is an LTR at each end, flanking an internal promoter; promoters useful in these vectors include those derived from chicken beta-actin, human histone, Herpes simplex thymidine kinase and rat thy 1.

10 Introduction of Genetic Material into Fibroblasts and
Assessment of Expression of the Genetic Material

A cell line which produces recombinant amphotropic retrovirus having a recombinant genome is used to infect fibroblasts. As described above, the recombinant genome can include a variety of components, but in general is comprised of two LTRs and, in place of the gag, the pol and the env sequences, a second promoter sequence and, in some cases, a gene encoding a selectable marker (e.g., neo).

20 Viral stocks, to be used in introducing genetic material of interest into fibroblasts, are harvested, as described above, supplemented with 8 micrograms per ml. (mcg/ml.) of Polybrene (Aldrich) and added to the culture of fibroblasts. If the titer of the virus is high (e.g., approximately 10^6 cfu per ml.), then virtually all fibroblasts will be infected and no selection (e.g., of fibroblasts into which the vector, including the recombinant genome, has been introduced) is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. If a

-23-

selectable marker is used, after exposure to the virus, the cells are grown to confluence and split into selective media (e.g., media containing G418 if the selectable marker is neo, media containing
5 histidinol and no histidine if the selectable marker is his).

The neo gene is a bacterial gene derived from the transposon Tn5, which encodes neomycin resistance in bacteria and resistance to the antibiotic G418 in
10 mammalian cells. This neo gene acts as a dominant selectable marker; its presence in a mammalian cell converts the cell into one which will grow in the presence of G418. (If it is not present, the cell dies in the presence of G418.) As a result, the
15 presence of this gene in a mammalian cell can be determined by culturing cells in media which contains G418. The recombinant retrovirus having this recombinant genome is referred to as the neo virus.

The recombinant retroviral vectors having the
20 neo gene also have a cloning site. As a result, genetic material of interest can be introduced into the vector, incorporated into fibroblasts and expressed by fibroblasts transduced with the recombinant retrovirus (referred to as fibroblasts
25 having incorporated genetic material).

At the BamHI cloning site, it is possible to insert genetic material of interest. The genetic material of interest can be DNA which does not normally occur in fibroblasts; DNA which normally
30 occurs in fibroblasts but is not expressed by them at levels which are biologically effective (i.e., levels sufficient to produce the normal physiological

-24-

effects of the polypeptide it encodes); DNA which occurs in fibroblasts and has been modified so that it is expressed by such cells; and any DNA which can be modified to be expressed by fibroblasts, alone or
5 in any combination thereof.

A copy of the gene encoding human parathyroid hormone (hPTH) has been cloned into this site, (e.g., into pLJ) in the following way: The pLJ plasmid was digested with BamHI and subsequently treated with the
10 enzyme calf intestinal phosphatase. Following this, the linear vector was fractionated on agarose gel and purified, using glass beads. In addition, the BamHI fragment containing the human PTH gene was prepared from the plasmid described by Hendy et al.,
15 which contains a complete cDNA of human PTH cloned into pBR322. Hendy, G.N. et al., Proceedings of the National Academy of Sciences, USA, 78:7365-7369 (1981).

A sub fragment of the PTH cDNA, containing 17 bp
20 of 5' untranslated, all coding and 31 bp of 3' untranslated sequence, was isolated by digesting the initial plasmid with DdeI and HinfI and isolating the 600bp fragment. The ends of this fragment were treated with DNA polymerase, to fill in the recessed
25 ends. BamHI linkers were ligated to the blunt ends with T_4 DNA ligase. An authentic BamHI restriction fragment was generated by digesting the ligation mixture from above with BamHI. This was then subcloned into the BamHI site of pBR322, which is the
30 plasmid used as the source of hPTH in vector construction.

-25-

Equal quantities of the pLJ linear backbone and the BamHI PTH fragment were added together, in the presence of T4 DNA ligase. The resulting mixture was maintained under conditions appropriate for ligation of the two fragments. The ligation mixture was used to transform bacteria HB101, which were then plated onto agar containing Kanamycin. Maniatis, T. et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, p.p. 250-251, 504; Bolivar, F. and K. Backman, In: Methods in Enzymology, R. Wu (ed.), Vol. 68, Academic Press, N.Y. (1979). The resulting colonies were analyzed for the recombinant plasmid.

Parathyroid hormone is a polypeptide which has a role in the regulation of calcium in the body. Although the hPTH gene is present in human fibroblasts, it is not expressed in those cells at biologically significant levels. Fibroblasts capable of making a polypeptide hormone such as hPTH, or another substance not normally made by such cells at biologically significant levels, can be engrafted onto an individual and serve as a continuous supply system for the hormone, or other substance.

This process is described with reference to hPTH, but it is to be understood that any gene can be introduced into fibroblasts, using the method described, and its expression by the cells can be assessed in a similar manner. As explained, DNA encoding the proteins or functions listed in Table 1 has been cloned into retroviral vectors and in many cases, introduced into fibroblasts in a similar manner and expressed by the transduced cells.

-26-

The Psi am cells producing the recombinant virus construct which contained the hPTH-encoding DNA and DNA encoding a selectable marker (such as the neo gene), were used to produce a viral stock, as described above. The viral stock was harvested; fibroblasts to be transduced with the virus containing the hPTH gene were incubated with the stock. In this case, a selectable marker is used to identify and select for transduced fibroblasts by culturing on media containing G418. If the viral titer is sufficiently high, that essentially all fibroblasts are infected and selection, using a selectable marker and appropriate media, is not needed.

Diploid fibroblasts from a Wistar rat transduced with DNA encoding hPTH according to the method of the present invention have been deposited at the American Type Culture Collection (Rockville, MD) under deposit number CRL9514. The ability of fibroblasts transduced with the recombinant retrovirus having the hPTH gene to express the hPTH gene has been assessed, both in vitro and in vivo.

In vitro assessment was carried out as follows:

The in vitro production of hPTH by NIH 3T3 cells infected with a virus containing the hPTH gene and selected in neomycin and by secondary cultures of fibroblasts established from a Wistar rat (FWR) was assessed. NIH 3T3 cells are cells derived from a chicken embryo and immortalized by selective cultivation in vitro. The FWR cells were derived from an explant from a syngeneic strain of rats

-27-

(Wistar). The cells have a limited life span and are referred to as secondary cultures.

Both types of cells had previously been infected with virus and selected in neomycin, as described
5 above. After selection, each type of cell was seeded onto 10 cm tissue culture dishes and grown to confluence. Fresh culture media (DME with 10% CS and penicillin and streptomycin) was then added; this point is referred to subsequently as time zero. At
10 subsequent times, aliquots of media were harvested and frozen. At the end of 24 hours all media was removed from the cell and placed back in the incubator; 12 hours later an additional aliquot was taken for analysis of the lability of the human PTH
15 in tissue culture.

The aliquots were analyzed for the presence of hPTH using a radioimmunoassay (Nichols) which measures intact hPTH. The technique is described in AllegroTM Intact PTH/Immunoassay System for the
20 Quantitative Determination of Human Intact Parathyroid Hormone in Serum, Nichols Institute Diagnostics, San Juan Capistrano, CA (36B-2170, Effective 7/86 Revised), the teachings of which are incorporated herein by reference. The assay has a
25 sensitivity of approximately one nanogram/milliliter serum (ng/ml) and was shown to be specific for human PTH in that it does not cross react with rat PTH. The results of the experiments are plotted as the production of hPTH, as measured by RIA, over time.
30 Results are shown in Figure 4.

The same analysis was also carried out after the Wistar rat fibroblasts which had been infected and

-28-

selected were seeded onto Cytodex beads. When the cells were subconfluent on the beads, fresh media was added; aliquots were taken and the production rate of PTH was measured. Under these conditions, the rate of production was twice that as measured in tissue culture on tissue culture plastic. In both cases, at the end of the assessment period, cells were harvested by trypsinization and counted.

In vivo assessment was conducted as follows:

Approximately 2×10^7 engineered fibroblasts were injected into the intraperitoneal cavity of syngeneic rats, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. Rats into which only phosphate buffer saline (PBS) or beads alone, in similar quantities, were injected intraperitoneally served as controls. A control rat received an injection intraperitoneally (IP) of PBS alone. A second control animal received an injection IP of 2×10^7 FWR cells expressing human PTH; they were administered in suspension in PBS. Another control rat received 300 mg of Cytodex 3 beads alone (i.e., without any cells). A fourth rat received approximately 150 mg of Cytodex 3 beads coated with FWR PTH cells, corresponding to approximately 2×10^7 cells. A fifth rat was administered approximately 200 mg of beads coated with FWR fibroblasts (approximately $2 - 3 \times 10^7$ cells). The time of injection is referred to as $T=0$.

Engraftment of the transduced fibroblasts was verified by determining the presence of hPTH in recipient's blood, using a radioimmunoassay specific

-29-

for hPTH. hPTH was detected in serum over a 3-week post-transplantation period.

In all animals, the first measurement was obtained at T=3 hours (3 hours after injection).

- 5 This was done by obtaining approximately 1/2 ml. of blood through the tail vein of each of the animals. Blood was then allowed to clot and the serum was analyzed for hPTH through the use of the Nichols radioimmunoassay described above. This was repeated;
- 10 at various times approximately 1/2 ml. of blood was obtained, through rat tail phlebotomy, and analyzed for hPTH. Results of these determinations are shown in the table. Also shown in the table are results of serum calcium levels in these animals.

TABLE 2
Assessment of in vivo expression of human PTH

TIME (DAYS)	RAT #1		RAT #2 ^a		RAT #3 ^b		RAT #4 ^c		RAT #5 ^c	
	Ca (mg/%)	hPTH (pg/ml.)	Ca (mg/%)	hPTH (pg/ml.)	Ca (mg/%)	hPTH (pg/ml.)	Ca (mg/%)	hPTH (pg/ml.)	Ca (mg/%)	hPTH (pg/ml.)
-3	-	-	-	-	10.70	1.5	-	-	-	-
-2	-	-	-	-	9.0	1	10.62	1	10.47	1
-1	-	-	-	-	9.13	1	10.14	1	9.81	1
0	9.73	1	9.96	55	10.11	1	10.99	1	-	-
3hr.	8.48	1	7.54	55	10.13	1	9.22	170	9.79	120
1	-	-	8.81	-	9.29	1	9.47	1	9.79	30
2	10.69	1	10.66	1	10.18	1	10.58	1	10.35	1
3	9.89	1	10.10	1	9.90	1	9.97	1	10.44	1
4	10.29	1	9.51	1	9.62	1	9.33	1	9.49	21
5	9.68	1	9.92	1	8.28	1	9.83	1	9.73	39
6	10.04	1	9.82	1	-	-	-	-	-	-
7	9.61	1	9.75	1	9.82	1	9.7	1	9.1	1
8	10.1	1	9.74	1	10.2	1	10.22	1	10.6	1
9	9.02	1	8.66	1	-	-	-	-	-	-
10	8.70	1	8.68	1	-	-	-	-	-	-
19	-	-	-	-	9.46	1	9.61	35	9.64	40
20	-	-	-	-	-	1	-	1	-	1

a Injected with free PTH-fibroblasts (IP)
b Injected with beads
c Injected with PTH fibro-beads (Cytodex 2 or 3)
 T = 0 = time of injection

SUBSTITUTE SHEET

-31-

Results of this assessment are represented in Figure 5. Within 3 hours after injection of transduced fibroblasts alone, 60 picograms (pg.) hPTH were detected per milliliter (ml.) of recipient's serum. (Normal human serum PTH levels range from 10-50 pg/ml.) Within 3 hours after injection of transduced fibroblasts confluent on beads, hPTH levels of 120 pg/ml. or 175 pg/ml. serum were detected. As is also shown in the table, rats injected with transduced fibroblasts containing the hPTH gene did not become hypercalcemic. This is unexpected because experiments in vitro suggest hPTH is active in rat tissue. One possible explanation is that counter regulatory mechanisms were activated (i.e., calcitonin).

As a result, it has been demonstrated that transduced fibroblasts secrete a polypeptide hormone which is normally not secreted at biologically significant levels by fibroblasts and, additionally, that they can secrete this hormone at physiological (or greater) concentrations. Transplantation of the fibroblasts secreting human parathyroid hormone (here, into the intraperitoneal cavity) demonstrated that the phenotypic change conferred on the fibroblasts by the retrovirus (i.e., secretion of hPTH) is detected systemically in the recipient. Administration of transduced fibroblasts, in a similar manner, to parathyroidectomized rats can further demonstrate the effectiveness of hPTH produced.

Thus, in one embodiment of the method of the present invention, fibroblasts are transduced, using

-32-

a retroviral vector in which the recombinant genome includes DNA of interest and, optionally, DNA encoding a dominant selectable marker. Transduced cells are isolated and grown to sufficient numbers. 5 They are introduced into a recipient by transplantation (e.g., into the intraperitoneal cavity, into the central nervous system). Alternatively, fibroblasts are similarly transduced, isolated, grown and then transplanted in conjunction 10 with or bound to a matrix. This approach is particularly useful when long-term (several weeks or longer) engraftment is desired. The matrix used can be, as described, a support such as microcarrier beads, particularly collagen-coated microcarrier beads. The 15 matrix can be virtually any solid matrix of suitable size. A skin graft, which contains transduced fibroblasts, can be applied.

Introduction of Genetic Material Encoding
Polypeptides Other Than hPTH

20 Genes encoding polypeptides other than hPTH can also be introduced into fibroblasts by means of the retroviral vector. For example, a gene encoding human growth hormone (HGH) or a gene encoding insulin can be introduced into fibroblasts. Such genes can be 25 introduced into fibroblasts, alone or in combination with a gene encoding a selectable marker, such as the neo gene. HGH is a polypeptide of about 29,000 Daltons which is normally secreted only by the hypothalamus. Insulin is a polypeptide which 30 regulates glucose levels in the bloodstream.

-33-

These genes, as well as others (e.g., some of those listed above), have been introduced into fibroblasts in the same manner as described above for the hPTH gene and the resulting transduced
5 fibroblasts can be grafted onto an appropriate site on the body.

Use of Other Dominant Selectable Markers in the
Introduction of Genetic Material Encoding
Polypeptides

10 It is also possible to use dominant selectable markers other than the neo gene to introduce genetic material into fibroblasts. For example, the His D gene can be used for this purpose. The His D gene is a bacterial gene from Salmonella and encodes
15 histidinol dehydrogenase, a polypeptide which converts histidinol to histidine. Histidine is an essential amino acid; histidinol is an alcohol analogue of histidine and can be converted to histidine under the proper metabolic conditions. If
20 cells are grown in media containing histidinol but lacking histidine, those cells having the His D gene can convert histidinol to histidine. Because histidine is essential to their function, those cells which have the His D gene (and thus can make
25 histidine) will survive and those lacking the gene will not.

A retrovirus vector having the His D gene has been used to infect keratinocytes. The keratinocytes containing His D gene were selected by growing these
30 cells in media lacking histidine but containing histidinol. As expected, keratinocytes having the

-34-

His D gene formed colonies and grew to confluence; those lacking the gene did not. In fact, such cells occurred at a much higher frequency than those in which the neo gene was included. These same
5 techniques are useful in selecting fibroblasts containing DNA of interest.

As a result of this work, it is also possible to use independent dominant selectable markers (e.g., the neo gene and the His D gene) to introduce new
10 genetic material into fibroblasts. In the case of polypeptides which have two different subunits, for example, separate dominant selectable markers can be used to introduce the genetic information encoding the two subunits. In addition, two or more dominant
15 selectable markers can be used in the case of polypeptides which need to be specifically cleaved or processed in order to become active (e.g., insulin and parathyroid hormone). A gene encoding the necessary processing enzyme can be introduced along
20 with the gene encoding the polypeptide hormone requiring such processing. This would enable fibroblasts to process that polypeptide hormone.

Other Vehicles for the Introduction of Genetic Material of Interest into Fibroblasts

25 It is also possible to use vehicles other than retroviruses to genetically engineer or modify fibroblasts. Genetic information of interest can be introduced into fibroblasts by means of any virus which can express the new genetic material in such
30 cells. For example, SV40, herpes virus, adenovirus

WO 89/02468

PCT/US88/03089

-35-

and human papilloma virus can be used for this purpose.

Introduction of Genetic Material of Interest into Other Types of Cells

5 Using the method of the present invention, it is also possible to incorporate genetic material of interest into other types of cells. The method can be used to introduce genetic material of interest into any system in which cells can be harvested from
10 tissue, grown in tissue culture on a solid matrix and infected with a retrovirus. Genetic material of interest can also be incorporated into the central nervous system. For example, it can be introduced into fibroblasts or into glial cells, or neuroglial
15 cells, which are non-neuronal cellular elements of the central and the peripheral nervous systems. The transduced cells can be introduced into the recipient animal. This can be done through stereotactic administration, directly into specific areas of the
20 brain, or by being introduced into the cerebrospinal fluid. In the central nervous system, glial cells include oligodendroglia cells, astrocytes, ependymal cells and microglia cells. In the peripheral nervous system, they include the satellite cells of ganglia
25 and the neurolemmal (or Schwann) cells around peripheral nerve fibers.

Genetic material of interest, such as the purine salvage enzyme hypoxanthine guanine phosphoribosyl transferase (HPRT) can be introduced into glial cells
30 for the purpose of correcting the inherited deficiency of HPRT which results in the Lesch-Nyhan

WO 89/02468

PCT/US88/03089

-36-

Syndrome; a devastating genetic disease with numerous CNS abnormalities. It can also be introduced into the central nervous system to treat or prevent, for example, degenerative diseases (e.g., by introducing genetic material encoding nerve growth factor), Parkinson's Disease (e.g., by introducing genetic material encoding DOPA synthetic enzymes) and genetic disorders caused by an "abnormal" gene (e.g., Huntington's Disease, manic depression, Alzheimer's Disease), in which the abnormal gene can be replaced, its function corrected or counteracted, or production of an encoded product altered.

Genetic material of interest can also be introduced into endothelial cells, which cover the serous membranes and line the heart, blood vessels, lymphatics and the anterior chamber of the eye. Genetic material of interest, such as hPTH, can be introduced into such cells in order to facilitate its delivery into the bloodstream.

20 Uses of Fibroblasts and Other Cells Having Incorporated Genetic Material

The present invention makes it possible to genetically engineer fibroblasts in such a manner that they produce polypeptides and proteins not normally produced in fibroblasts in biologically significant amounts and secrete them into the bloodstream or other area of the body, such as the central nervous system. The fibroblasts formed in this way can serve as a continuous drug delivery system to replace present regimens, which require

WO 89/02468

PCT/US88/03089

-37-

periodic administration (by ingestion, injection, etc.) of the needed substance.

For example, it can be used to provide continuous delivery of insulin, which at the present
5 time, must be isolated from the pancreas, extensively purified and then injected into the body by those whose insulin production or utilization is impaired. In this way, insulin can be introduced into the body via a continuous drug delivery system and, as a
10 result, there would be no need for daily injections of insulin.

Genetically engineered fibroblasts can also be used for the production of clotting factors. Hemophiliacs lack a protein called Factor VIII, which
15 is involved in clotting. Factor VIII is now administered by injection. Fibroblasts having genes encoding Factor VIII, can be used to make a skin graft in which they produce Factor VIII; as a skin graft, the tissue would secrete the factor into the
20 bloodstream.

Incorporation of genetic material of interest into fibroblasts and other types of cells would be particularly valuable in the treatment of inherited disease and the treatment of acquired disease. In
25 the case of inherited diseases, this approach is used to provide genetically modified fibroblasts and other cells which can be used as a metabolic sink. That is, such fibroblasts would serve to degrade a potentially toxic substance. For example, this could
30 be used in treating urea cycle disorders.

Fibroblasts of the present invention can also be used in the treatment of genetic diseases in which a

-38-

product (e.g., an enzyme or hormone) normally produced by the body is not produced or is made in insufficient quantities. Here, fibroblasts transduced with a gene encoding the missing or inadequately produced substance can be used to produce it in sufficient quantities. For example, this can be used in producing alpha-1 antitrypsin. It can also be used in the production of Factor VIII and Factor IX and thus would be useful in treating hemophilia.

There are many acquired diseases for which treatment can be provided through use of genetically engineered fibroblasts (i.e., fibroblasts transduced with genetic material of interest). For example, such cells can be used in treating anemia, which is commonly present in chronic disease and often associated with chronic renal failure (e.g., in hemodialysis patients). In this case, fibroblasts having incorporated in them a gene encoding erythropoietin would correct the anemia by stimulating the bone marrow to increase erythropoiesis (i.e. production of red blood cells).

Fibroblasts of the present invention can also be used to administer a low dose of tissue plasminogen activator as an activator to prevent the formation of thrombi. In this case, fibroblasts having incorporated genetic material which encodes TPA would be transplanted into an individual in whom thrombus prevention is desired. This would be useful, for example, as a prophylactic against common disorders such as coronary artery disease, cerebrovascular disease, peripheral vascular occlusive disease, vein

WO 89/02468

PCT/US88/03089

-39-

(e.g., superficial) thrombosis, such as seen in pulmonary emboli, or deep vein thrombosis.

Fibroblasts which contain DNA encoding calcitonin can be used in the treatment of Paget's Disease, a
5 progressive, chronic disorder of bone metabolism, in which calcitonin is presently administered subcutaneously.

Another application for a skin graft having genetically engineered fibroblasts is in birth
10 control. Tests are underway now for using a polypeptide hormone called lutenizing hormone releasing hormone (LHRH) in regulating fertility. Continuous administration of LHRH results in a sterile individual; when administration ceases, the
15 individual is again fertile. Rather than taking LHRH injections or oral medication, one could have a small graft which continuously secretes LHRH to provide the same effect. In the event that the person wanted to regain fertility this transplant could be excised;
20 delivery of the polypeptide hormone would cease.

Fibroblasts engineered to produce and secrete interleukins (e.g., IL-1, IL-2, IL-3) can be used in several contexts. For example, the result of some of the therapies now used (e.g., chemotherapy) is
25 induction of neutropenia (the presence of abnormally low numbers of neutrophils in the blood), often caused by direct suppression of the bone marrow. For example, use of virtually all the chemotherapeutic agents, as well as AZT, used in the treatment of
30 (AIDS) Acquired Immune Deficiency Syndrome, results in neutropenia. This condition results in numerous life-threatening infections. In these cases,

WO 89/02468

PCT/US88/03089

-40-

administration of, for example, IL-3 through
transplantation of fibroblasts which contain genetic
material encoding IL-3 can be used to increase the
neutrophil count. In addition, the administration of
5 thrombopoietin, which stimulates the production of
platelets, can be used in the treatment of numerous
conditions in which platelet count is low. In this
case, fibroblasts transduced with the gene for
thrombopoietin can be applied to an individual;
10 production and secretion of the encoded product will
result in stimulation of platelet production.

Another related application of fibroblasts
having incorporated genetic material is in the
treatment of AIDS. Interleukin 2 and Interleukin 3,
15 which stimulate the immune system, are potentially
valuable in the treatment of AIDS. They could be
delivered by a skin graft having fibroblasts which
have been genetically engineered to produce these two
polypeptides (which are now administered by periodic
20 injection).

Another use of the present invention is in the
treatment of enzyme defect diseases. In this case
the product (polypeptide) encoded by the gene
introduced into fibroblasts is not secreted (as are
25 hormones); rather, it is an enzyme which remains
inside the cell. There are numerous cases of genetic
diseases in which an individual lacks a particular
enzyme and is not able to metabolize various amino
acids or other metabolites. The correct genes for
30 these enzymes could be introduced into a skin
transplant; the transplant would then carry out that
metabolic function. For example, there is a genetic

WO 89/02468

PCT/US88/03089

-41-

disease in which those affected lack the enzyme adenosine deaminase. This enzyme is involved in the degradation of purines to uric acid. It might be possible, using the present invention, to produce a
5 skin graft capable of producing the missing enzyme at sufficiently high levels to detoxify the blood as it passes through the area to which the graft is applied.

The present invention also has veterinary
10 applications. It can be used, for example, in delivering substances such as drugs (e.g., antibiotics) and hormones to animals, which would otherwise be provided by being incorporated into their feed, added to their water or injected
15 periodically (e.g., daily or less frequently). Use of the modified fibroblasts of the present invention has the advantage that the tissue formed of the modified cells can be applied to the animal and will provide quantities of the encoded protein on an
20 ongoing basis, thus eliminating the need for daily/periodic administration of the substance.

WO 89/02468

PCT/US88/03089

-42-

Industrial Utility

This invention has industrial applicability in providing hormones, enzymes and drugs to mammals, including humans, in need of such substances. For example, it can be used to provide a continuous supply of a hormone which otherwise would be administered on a intravenously, intramuscularly or subcutaneously. It is particularly valuable in providing such substances, such as hormones (e.g., parathyroid hormone, insulin), which are needed for extended periods of time.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described specifically herein. Such equivalents are intended to be encompassed in the scope of the following claims.

-43-

CLAIMS

1. Fibroblasts capable of expressing incorporated genetic material which is not normally expressed at biologically significant levels in
5 fibroblasts.
2. Fibroblasts of Claim 1 in which the incorporated genetic material is DNA or RNA which does not normally occur in fibroblasts; DNA or RNA which normally occurs in fibroblasts but is not
10 expressed in them at levels which are biologically significant; DNA or RNA which normally occurs in fibroblasts and has been modified so that it is expressed in fibroblasts; and any DNA or RNA which can be modified to be
15 expressed by fibroblasts, alone or in any combination thereof.
3. Fibroblasts of Claim 2, wherein the incorporated genetic material additionally comprises genetic material encoding at least one selectable
20 marker.
4. Fibroblasts of Claim 3 in which at least one selectable marker is a dominant selectable marker.
5. Fibroblasts of Claim 4 in which the dominant
25 selectable marker is a gene encoding antibiotic resistance.

WO 89/02468

PCT/US88/03089

-44-

6. Human fibroblasts capable of expressing incorporated genetic material not normally expressed by human fibroblasts at biologically significant levels.
- 5 7. Human fibroblasts of Claim 6 in which the incorporated genetic material encodes a hormone, an enzyme or a drug.
8. Human fibroblasts of Claim 7, wherein the incorporated genetic material additionally
10 comprises genetic material which encodes at least one dominant selectable marker.
9. Fibroblasts containing incorporated genetic material encoding a) a polypeptide not normally made at biologically significant levels in
15 fibroblasts and a b) neo gene, said neo gene being a bacterial gene derived from the transposon Tn5.
10. Human fibroblasts capable of expressing a polypeptide not normally made at biologically
20 significant levels in human fibroblasts.
11. Human fibroblasts expressing incorporated genetic material, the incorporated genetic material being selected from the group consisting of: a) genetic material encoding
25 hormone not normally made at biologically significant levels in fibroblasts; b) genetic material encoding an enzym not normally made at

WO 89/02468

PCT/US88/03089

-45-

- biologically significant levels in fibroblasts;
c) genetic material encoding a drug not normally
made at biologically significant levels in
fibroblasts; and d) genetic material encoding at
least one dominant selectable marker.
12. Fibroblasts having incorporated therein
recombinant amphotropic retrovirus having a
recombinant genome comprised of:
- a. long terminal repeat sequences, the tRNA
binding site and the Psi packing site derived
from amphotropic Moloney murine leukemia
virus; and
- b. genetic material encoding at least one
polypeptide of interest.
13. Fibroblasts of Claim 13 in which the recombinant
genome additionally comprises genetic material
encoding at least one selectable marker.
14. Fibroblasts of Claim 13 in which the genetic
material encoding at least one selectable marker
is a neo gene.
15. Fibroblasts of Claim 14 in which the genetic
material encodes human parathyroid hormone.
16. A method of making fibroblasts which express
genetic material of interest which encodes at
least one polypeptide of interest, comprising:
- a. contacting cultured fibroblasts with media
containing an infectious recombinant retrovirus

WO 89/02468

PCT/US88/03089

-46-

- having a recombinant genome comprised of incorporated genetic material; and
- b. maintaining the cultured fibroblasts and the media containing the infectious recombinant retrovirus under conditions appropriate for infection of the fibroblasts by recombinant retrovirus.
17. A method of Claim 16 in which the genetic material of interest encodes a hormone, an enzyme or a drug not normally expressed in fibroblasts at biologically significant levels.
18. A method of Claim 17 in which the recombinant genome of the retrovirus additionally comprises genetic material encoding at least one dominant selectable marker.
19. A method of making fibroblasts, capable of expressing incorporated genetic material encoding at least one protein or at least one polypeptide of interest, comprising:
- a. transducing cultured fibroblasts with genetic material encoding at least one protein or at least one polypeptide of interest and
- b. culturing the transduced fibroblasts under conditions appropriate for their growth.
20. A method of providing a hormone, an enzyme or a drug to the body comprising transducing fibroblasts with genetic material encoding the hormone, the enzyme or the drug and introducing

WO 89/02468

PCT/US88/03089

-47-

the transduced fibroblasts into the body or applying the transduced fibroblasts to the surface of the body.

21. A method of providing a hormone, an enzyme or a drug to the body, comprising:
- 5 a. incorporating into fibroblasts a recombinant retrovirus, the recombinant retrovirus having a recombinant genome comprised of:
- 10 1) genetic material encoding the hormone, the enzyme or the drug;
- 15 2) the long terminal repeat sequences, the tRNA binding site and the Psi packaging site derived from an amphotropic retrovirus; and
- 20 3) at least one promoter of eukaryotic origin; and
- b. introducing the resulting fibroblasts into the body or applying the resulting fibroblasts to the surface of the body.
22. A method of Claim 21 in which the recombinant genome is additionally comprised of a promoter of eukaryotic origin which can be modulated by an external cue.
- 25 23. A method of Claim 22 in which the recombinant genome is additionally comprised of genetic material encoding at least one dominant selectable marker.

WO 89/02468

PCT/US88/03089



-48-

24. A method of Claim 23 in which at least one dominant selectable marker is a gene encoding antibiotic resistance or a gene that complements a genetic defect in the host species.
- 5 25. A method of controlling fertility, comprising:
- 10 a. forming a transplantable sheet of fibroblasts expressing at biologically significant levels at least one hormone which regulates fertility; and
- b. applying said transplantable sheet to the body.

WO 89/02468

1 / 4

PCT/US88/03089



*Fig. 1**Fig. 4*

WO 89/02468

PCT/US88/03089


2 / 4

VECTORS WITH SELECTABLE MARKERS



VECTORS WITHOUT SELECTABLE MARKERS



*Fig. 3*

WO 89/02468

PCT/US88/03089

4 / 4




Fig. 5

INTERNATIONAL SEARCH REPORT

International Application No PCT/US88/03089

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC ⁴ : C 12 N 15/00		
II. FIELDS SEARCHED		
Minimum Documentation Searched *		
Classification System	Classification Symbols	
IPC ⁴	C 12 N	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched *		
III. DOCUMENTS CONSIDERED TO BE RELEVANT *		
Category *	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X Y	Proc. Natl. Acad. Sci. Vol. 81, October 1984, R D Cone et al.: "High-efficiency gene transfer into mammalian cells: Generation of helper free recombinant retrovirus with broad mammalian host range", pages 6349-6353, see the whole article --	1-6 7-19
X	Proc. Natl. Acad. Sci. Vol. 84, February 1987, I D Palmer et al.: "Efficient retrovirus-mediated transfer and expression of a human adenosine deaminase gene in diploid skin fibroblasts from an adenosine-deficient human", pages 1055-1059 see the whole article --	1-19
X	Proc. Natl. Acad. Sci. Vol. 84, February 1987, J Sorge et al.: "Complete correction of enzymatic defect of type I: Gaucher disease fibroblasts by retroviral-mediated gene transfer", pages 906-909. see the whole article --	1-19 ./.
<p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
8th December 1988	12 JAN 1989	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	P.C.G. VAN DER PUTTEN	

International Application No. PCT/US88/03089

III. D CUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
X Y	Journal of Virology, Vol. 50(2), May 1984, L-H Sherry Hwang et al.: "Expression of genes introduced into cells by retroviral infection is more efficient than that of genes introduced into cells by DNA transfection", pages 417-424, see the whole article --	1-5, 9 6-8, 12-14, 16, 18
X Y	Proc. Natl. Acad. Sci., Vol. 79, April 1982, J Doeimer et al.: "Introduction of rat growth hormone gene into mouse fibroblasts via a retroviral DNA vector: Expression and regulation", pages 2268-2272 see the whole article --	1-5, 9 12-19
	Molecular and Cell. Biol. Vol. 7(8), August 1987, T G Hugo et al.: "Isolation of microcell hybrid clones containing retroviral vector insertions into specific human chromosomes", pages 2814-2820 see the whole article --	1-6 7-19
X	Journal of Virology, Vol. 61(8), August 1987, B Seliger et al.: "Murine gamma interferon inhibits v-mos-induced fibroblast transformation via down regulation of retroviral gene expression", pages 2567-2572 see the whole article --	1-5 9
X	EP, A, 0 178 220 (INSTITUT PASTEUR) 16 April 1984 see the whole document --	1-19
X Y	US, A, 4 634 665 (RAXEL et al.) 6 January 1987 see the whole document -----	1-5 6-11

International Application No. PCT/US88/03089

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹

This International search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers 20-25 because they relate to subject matter not required to be searched by this Authority, namely:

Methods for treatment of the human or animal body; see rule 39.1 (IV).

2. ☐ Claim numbers..... because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers..... because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ²

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

PCT/US88/03089
SA 24341

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EIMP file on 02/11/88. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0178220	16-04-86	None	
US-A- 4634665	06-01-87	WO-A- 8102426	03-09-81
		EP-A- 0045809	17-02-82
		US-A- 4399216	16-08-83
		CA-A- 1179953	25-12-84
		AU-A- 558061	15-01-87